### **GYNAECOMASTIA**



### **DEVELOPMENT OF BREAST TISSUE IN MEN**

AN ENLARGEMENT OF 0.5 CM OF BREAST TISSUE

### **CLASSIFICATIONS**

• 1<sup>ST</sup> CLASSIFICATION-

**TRUE** 

**PSEUDO** 

. 2<sup>ND</sup> CLASSIFICATION-

CONGENITAL ACQUIRED

- 1. PHYSIOLOGIC
- 2. PATHOLOGIC

## SIMON CLASSIFICATION




GRADE ENLARGEMENT SKIN LAXITY

• 1 SMALL ABSENT

2A MODERATE ABSENT

2B MODERATE PRESENT

• 3 LARGE PRESENT



### CONGENITAL GYNAECOMASTIA

#### **PHYSIOLOGIC:**

NEONATAL PUBERTAL SENILE

#### **PATHOLOGIC:**

HYPOGONADAL ANDOGEN RESISTANCE

ENZYMATIC DEFECT OF TESTOSTERONE SYNTHESIS

CONGENITAL ANORCHISM


KLINEFELTER SYNDROME

HYPERESTOGENIC TRUE HERMAPHRODITISM

CONGENITAL ADRENAL HYPERPLASIA ELEVATED PERIPHERAL AROMATASE

# INFERIOR SEMICIRCULAR AREOLA INCISIA





 PERIAREOLAR INCISION PREFERABLY BE WAVY RATHER THAN SEMICIRCULAR TO PREVENT FROM BEING NOTICED AND AVOID AN OPERATED LOOK.

( AS DESCRIBED BY DR. WOFFLES WU, SINGAPORE stealth incision )

### **GRADE 1**



- SMALL PROTUBERANCE OF FIBROUS TISSUE BELOW THE AREOLA.
- SIMPLE EXCISION THROUGH SMALL PERIAREOLA OR TRANSEVERE INCISION.
- 29% NIPPLE RETRACTION
- S A L OR U A L IS THE TREATMENT AS AN ADJUNCT TO EXCISION

### **GRADE 2**



- EXCISION AND S A L IS THE TREATMENT OPTION.
- S A L IS SATISFACTORY TREATMENT FOR LARGER BREAST WITHOUT PTOSIS AND WITHOUT EXCESSIVE FIBROSIS.

### **GRADE 3**



- PTOSIS IS NOT SEVERE---SUCTION
  LIPECTEOMY AND REPEAT AFTER 3MONTHS.
- PTOSIS IS SEVERE--- SKIN REDUCTION IN COMBINATION WITH SUB CUTANEOUS MASTECTOMY WITH NIPPLE GRAFTING.
   (INFRA MAMMARY INCISION)

### **TUMESCENT SOLUTION FOR S A L**

- NORMAL SALINE 100ML
- 2 AMPS 1 IN 1000ADRENALINE
- 2% XYLOCAINE 20ML
- 0.5% BUPIVACAINE 5ML
- SODABICARB 10ML
- HYALURONIDASE 1500 I U
- TRIAMCINOLONE 10 mg/ml 1 ml.





